

# AIEEE- 2004 Chemistry Paper

76. Which of the following sets of quantum numbers is correct for an electron in 4f orbital?

(1)  $n = 4, l = 3, m = +4, s = +\frac{1}{2}$                       (2)  $n = 3, l = 2, m = -2, s = +\frac{1}{2}$

(3)  $n = 4, l = 3, m = +1, s = +\frac{1}{2}$                       (4)  $n = 4, l = 4, m = -4, s = -\frac{1}{2}$

**Ans.**  $n = 4, l = 3, m = +1, s = +\frac{1}{2}$

77. Consider the ground state of Cr atom ( $Z = 24$ ). The number of electrons with the azimuthal quantum numbers  $l = 1$  and  $2$  are respectively

(1) 12 and 4    (2) 16 and 5

(3) 16 and 4    (4) 12 and 5

**Ans.** 12 and 5

78. Which one of the following ions has the highest value of ionic radius?

(1)  $\text{Li}^+$     (2)  $\text{F}^-$

(3)  $\text{O}^{2-}$     (4)  $\text{B}^{3+}$

**Ans.**  $\text{O}^{2-}$

79. The wavelength of the radiation emitted, when in hydrogen atom electron falls from infinity to stationary state 1, would be (Rydberg constant =  $1.097 \times 10^7 \text{ m}^{-1}$ )

(1) 91 nm    (2)  $9.1 \times 10^{-8}$  nm

(3) 406 nm    (4) 192 nm

**Ans.** 91 nm

80. The correct order of bond angles (smallest first) in  $\text{H}_2\text{S}$ ,  $\text{NH}_3$ ,  $\text{BF}_3$  and  $\text{SiH}_4$  is

(1)  $\text{H}_2\text{S} < \text{SiH}_4 < \text{NH}_3 < \text{BF}_3$                       (2)  $\text{H}_2\text{S} < \text{NH}_3 < \text{BF}_3 < \text{SiH}_4$

(3)  $\text{H}_2\text{S} < \text{NH}_3 < \text{SiH}_4 < \text{BF}_3$                       (4)  $\text{NH}_3 < \text{H}_2\text{S} < \text{SiH}_4 < \text{BF}_3$

**Ans.**  $\text{H}_2\text{S} < \text{NH}_3 < \text{SiH}_4 < \text{BF}_3$

81. Which one of the following sets of ions represents the collection of isoelectronic species?

(1)  $\text{K}^+, \text{Ca}^{2+}, \text{Sc}^{3+}, \text{Cl}^-$                               (2)  $\text{Na}^+, \text{Mg}^{2+}, \text{Al}^{3+}, \text{Cl}^-$

(3)  $\text{K}^+, \text{Cl}^-, \text{Mg}^{2+}, \text{Sc}^{3+}$                               (4)  $\text{Na}^+, \text{Ca}^{2+}, \text{Sc}^{3+}, \text{F}^-$

**Ans.**  $\text{K}^+, \text{Ca}^{2+}, \text{Sc}^{3+}, \text{Cl}^-$

82. Among  $\text{Al}_2\text{O}_3$ ,  $\text{SiO}_2$ ,  $\text{P}_2\text{O}_3$  and  $\text{SO}_2$  the correct order of acid strength is

(1)  $\text{SO}_2 < \text{P}_2\text{O}_3 < \text{SiO}_2 < \text{Al}_2\text{O}_3$                       (2)  $\text{Al}_2\text{O}_3 < \text{SiO}_2 < \text{P}_2\text{O}_3 < \text{SO}_2$

(3)  $\text{Al}_2\text{O}_3 < \text{SiO}_2 < \text{SO}_2 < \text{P}_2\text{O}_3$                       (4)  $\text{SiO}_2 < \text{SO}_2 < \text{Al}_2\text{O}_3 < \text{P}_2\text{O}_3$

**Ans.**  $\text{Al}_2\text{O}_3 < \text{SiO}_2 < \text{P}_2\text{O}_3 < \text{SO}_2$

83. The bond order in  $\text{NO}$  is 2.5 while that in  $\text{NO}^+$  is 3. Which of the following statements is true for these two species?

(1) Bond length in  $\text{NO}^+$  is greater than in  $\text{NO}$


(2) Bond length is unpredictable

(3) Bond length in  $\text{NO}^+$  is equal to that in  $\text{NO}$

(4) Bond length in  $\text{NO}$  is greater than in  $\text{NO}^+$

**Ans.** Bond length in NO is greater than in NO<sup>+</sup>

84. The formation of the oxide ion O<sup>2-</sup>(g) requires first an exothermic and then an endothermic step as shown below



- (1) Oxygen is more electronegative
- (2) O<sup>-</sup> ion has comparatively larger size than oxygen atom
- (3) O<sup>-</sup> ion will tend to resist the addition of another electron
- (4) Oxygen has high electron affinity

**Ans.** O<sup>-</sup> ion will tend to resist the addition of another electron

85. The states of hybridization of boron and oxygen atoms in boric acid (H<sub>3</sub>BO<sub>3</sub>) are respectively

- (1) sp<sup>2</sup> and sp<sup>2</sup>
- (2) sp<sup>3</sup> and sp<sup>3</sup>
- (3) sp<sup>3</sup> and sp<sup>2</sup>
- (4) sp<sup>2</sup> and sp<sup>3</sup>

**Ans.** sp<sup>2</sup> and sp<sup>3</sup>

86. Which one of the following has the regular tetrahedral structure?

- (1) XeF<sub>4</sub>
- (2) [Ni(CN)<sub>4</sub>]<sup>2-</sup>
- (3) BF<sub>4</sub><sup>-</sup>
- (4) SF<sub>4</sub>

**Ans.** BF<sub>4</sub><sup>-</sup>

87. Of the following outer electronic configurations of atoms, the highest oxidation state is achieved by which one of them?

- (1) (n-1)d<sup>8</sup>ns<sup>2</sup>
- (2) (n-1)d<sup>5</sup>ns<sup>2</sup>
- (3) (n-1)d<sup>3</sup>ns<sup>2</sup>
- (4) (n-1)d<sup>5</sup>ns<sup>-1</sup>

**Ans.** (n-1)d<sup>5</sup>ns<sup>2</sup>

88. As the temperature is raised from 20°C to 40°C, the average kinetic energy of neon atoms changes by a factor of which of the following?

- (1) ½
- (2) 2
- (3)  $\frac{313}{293}$
- (4)  $\sqrt{\frac{313}{293}}$

**Ans.**  $\frac{313}{293}$

89. The maximum number of 90° angles between bond pair of electrons is observed in

- (1) dsp<sup>3</sup> hybridization
- (2) sp<sup>3</sup>d<sup>2</sup> hybridization
- (3) dsp<sup>2</sup> hybridization
- (4) sp<sup>3</sup>d hybridization

**Ans.** sp<sup>3</sup>d<sup>2</sup> hybridization

90. Which one of the following aqueous solutions will exhibit highest boiling point?

- (1) 0.01 M Na<sub>2</sub>SO<sub>4</sub>
- (2) 0.015 M glucose
- (3) 0.015 M urea
- (4) 0.01 M KNO<sub>3</sub>

**Ans.** 0.01 M Na<sub>2</sub>SO<sub>4</sub>

91. Which among the following factors is the most important in making fluorine the strongest oxidizing halogen?

- (1) Electron affinity  
(2) Bond dissociation energy  
(3) Hydration enthalpy  
(4) Ionization enthalpy

**Ans.** Bond dissociation energy

92. In Vander Waals equation of state of the gas law, the constant 'b' is a measure of  
(1) intermolecular repulsions  
(2) intermolecular collisions per unit volume  
(3) Volume occupied by the molecules  
(4) intermolecular attraction

**Ans.** Volume occupied by the molecules

93. The conjugate base of  $\text{H}_2\text{PO}_4^-$  is  
(1)  $\text{PO}_4^{3-}$   
(2)  $\text{HPO}_4^{2-}$   
(3)  $\text{H}_3\text{PO}_4$   
(4)  $\text{P}_2\text{O}_5$

**Ans.**  $\text{HPO}_4^{2-}$

94.  $6.02 \times 10^{20}$  molecules of urea are present in 100 ml of its solution. The concentration of urea solution is  
(1) 0.001 M  
(2) 0.1 M  
(3) 0.02 M  
(4) 0.01 M

**Ans.** 0.01 M

95. To neutralize completely 20 mL of 0.1 M aqueous solution of phosphorous acid ( $\text{H}_3\text{PO}_3$ ), the volume of 0.1 M aqueous KOH solution required is  
(1) 10 mL  
(2) 60 mL  
(3) 40 mL  
(4) 20 mL

**Ans.** 40 mL

96. For which of the following parameters the structural isomers  $\text{C}_2\text{H}_5\text{OH}$  and  $\text{CH}_3\text{OCH}_3$  would be expected to have the same values?  
(Assume ideal behaviour)  
(1) Heat of vaporization  
(2) Gaseous densities at the same temperature and pressure  
(3) Boiling points  
(4) Vapour pressure at the same temperature

**Ans.** Gaseous densities at the same temperature and pressure


97. Which of the following liquid pairs shows a positive deviation from Raoult's law?  
(1) Water – hydrochloric acid  
(2) Acetone – chloroform  
(3) Water – nitric acid  
(4) Benzene – methanol

**Ans.** Benzene – methanol

98. Which one of the following statements is false?  
(1) Raoult's law states that the vapour pressure of a components over a solution is proportional to its mole fraction  
(2) Two sucrose solutions of same molality prepared in different solvents will have the same freezing point depression  
(3) The correct order of osmotic pressure for 0.01 M aqueous solution of each compound is  $\text{BaCl}_2 > \text{KCl} > \text{CH}_3\text{COOH} > \text{sucrose}$   
(4) The osmotic pressure ( $\pi$ ) = MRT, where M is the molarity of the solution

**Ans.** Two sucrose solutions of same molality prepared in different solvents will have the same freezing point depression

99. What type of crystal defect is indicated in the diagram below?



- (1) Frenkel defect (2) Frenkel and Schottky defects  
(3) Interstitial defect (4) Schottky defect

**Ans.** Schottky defect

100. An ideal gas expands in volume from  $1 \times 10^{-3} \text{ m}^3$  to  $1 \times 10^{-2} \text{ m}^3$  at 300 K against a constant pressure of  $1 \times 10^5 \text{ Nm}^{-2}$ . The work done is

- (1) -900 J (2) 900 kJ  
(3) 2780 kJ (4) -900 kJ

**Ans.** -900 J

101. In hydrogen – oxygen fuel cell, combustion of hydrogen occurs to

- (1) generate heat  
(2) remove adsorbed oxygen from electrode surfaces  
(3) produce high purity water  
(4) create potential difference between the two electrodes

**Ans.** create potential difference between the two electrodes

102. In first order reaction, the concentration of the reactant decreases from 0.8 M to 0.4 M in 15 minutes. The time taken for the concentration to change from 0.1 M to 0.025 M is

- (1) 30 minutes (2) 60 minutes  
(3) 7.5 minutes (4) 15 minutes

**Ans.** 30 minutes

103. What is the equilibrium expression for the reaction  $\text{P}_{4(s)} + 5\text{O}_{2(g)} \rightleftharpoons \text{P}_4\text{O}_{10(s)}$ ?

- (1)  $K_c = [\text{P}_4\text{O}_{10}] / [\text{P}_4][\text{O}_2]^5$  (2)  $K_c = 1/[\text{O}_2]^5$   
(3)  $K_c = [\text{O}_2]^5$  (4)  $K_c = [\text{P}_4\text{O}_{10}] / 5[\text{P}_4][\text{O}_2]$

**Ans.**  $K_c = 1/[\text{O}_2]^5$

104. For the reaction,  $\text{CO}(g) + \text{Cl}_2(g) \rightleftharpoons \text{COCl}_2(g)$  the  $\frac{K_p}{K_c}$  is equal to

- (1)  $\frac{1}{RT}$  (2) 1.0  
(3)  $\sqrt{RT}$  (4) RT

**Ans.**  $\frac{1}{RT}$

105. The equilibrium constant for the reaction  $\text{N}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}(g)$  at temperature T is  $4 \times 10^{-4}$ . The value of  $K_c$  for the reaction  $\text{NO}(g) \rightleftharpoons \frac{1}{2}\text{N}_2(g) + \frac{1}{2}\text{O}_2(g)$  at the same temperature is

- (1)  $2.5 \times 10^2$  (2) 0.02  
 (3)  $4 \times 10^{-4}$  (4) 50

**Ans.** 50

106. The rate equation for the reaction  $2A + B \longrightarrow C$  is found to be: rate  $k[A][B]$ . The correct statement in relation to this reaction is that the

- (1) unit of  $K$  must be  $s^{-1}$   
 (2) values of  $k$  is independent of the initial concentration of A and B  
 (3) rate of formation of C is twice the rate of disappearance of A  
 (4)  $t_{1/2}$  is a constant


**Ans.** values of  $k$  is independent of the initial concentration of A and B

107. Consider the following  $E^\circ$  values

$$E^\circ_{\text{Fe}^{3+}/\text{Fe}^{2+}} = 0.77 \text{ V}$$

$$E^\circ_{\text{Sn}^{2+}/\text{Sn}} = -0.14 \text{ V}$$

Under standard conditions the potential for the reaction



- (1) 1.68 V (2) 0.63 V  
 (3) 0.91 V (4) 1.40 V

**Ans.** 0.91 V

108. The molar solubility product is  $K_{sp}$ . 's' is given in terms of  $K_{sp}$  by the relation

- (1)  $s = \left(\frac{K_{sp}}{128}\right)^{1/4}$  (2)  $s = \left(\frac{K_{sp}}{256}\right)^{1/5}$   
 (3)  $s = (256K_{sp})^{1/5}$  (4)  $s = (128K_{sp})^{1/4}$

**Ans.**  $s = \left(\frac{K_{sp}}{256}\right)^{1/5}$

109. The standard e.m.f of a cell, involving one electron change is found to be 0.591 V at  $25^\circ\text{C}$ . The equilibrium constant of the reaction is ( $F = 96,500 \text{ C mol}^{-1}$ ;  $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ )

- (1)  $1.0 \times 10^1$  (2)  $1.0 \times 10^{30}$   
 (3)  $1.0 \times 10^{10}$  (4)  $1.0 \times 10^5$

**Ans.**  $1.0 \times 10^{10}$

110. The enthalpies of combustion of carbon and carbon monoxide are  $-393.5$  and  $-283 \text{ kJ mol}^{-1}$  respectively. The enthalpy of formation of carbon monoxide per mole is

- (1) 110.5 kJ (2) -110.5 kJ  
 (3) -676.5 kJ (4) 676.5 kJ

**Ans.** -110.5 kJ

111. The limiting molar conductivities  $\Lambda^\circ$  for NaCl, KBr and KCl are 126, 152 and  $150 \text{ S cm}^2 \text{ mol}^{-1}$  respectively. The  $\Lambda^\circ$  for NaBr is

- (1)  $128 \text{ S cm}^2 \text{ mol}^{-1}$  (2)  $302 \text{ S cm}^2 \text{ mol}^{-1}$   
 (3)  $278 \text{ S cm}^2 \text{ mol}^{-1}$  (4)  $176 \text{ S cm}^2 \text{ mol}^{-1}$

**Ans.**  $128 \text{ S cm}^2 \text{ mol}^{-1}$

112. In a cell that utilises the reaction  $\text{Zn(s)} + 2\text{H}^+(\text{aq}) \longrightarrow \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$  addition of  $\text{H}_2\text{SO}_4$  to cathode compartment, will
- (1) lower the E and shift equilibrium to the left
  - (2) increases the E and shift equilibrium to the left
  - (3) increase the E and shift equilibrium to the right
  - (4) Lower the E and shift equilibrium to the right

**Ans.** increase the E and shift equilibrium to the right

113. Which one the following statement regarding helium is incorrect?
- (1) It is used to fill gas balloons instead of hydrogen because it is lighter and non – inflammable
  - (2) It is used in gas – cooled nuclear reactors
  - (3) It is used to produce and sustain powerful superconducting reagents
  - (4) It is used as cryogenic agent for carrying out experiments at low temperatures

**Ans.** It is used to fill gas balloons instead of hydrogen because it is lighter and non – inflammable

114. Identify the correct statements regarding enzymes
- (1) Enzymes are specific biological catalysts that can normally function at very high temperature ( $T \sim 1000 \text{ K}$ )
  - (2) Enzymes are specific biological catalysts that the posses well – defined active sites
  - (3) Enzymes are specific biological catalysts that can not be poisoned
  - (4) Enzymes are normally heterogeneous catalysts that are very specific in their action

**Ans.** Enzymes are specific biological catalysts that the posses well – defined active sites

115. One mole of magnesium nitride on the reaction with an excess of water gives
- |                          |                              |
|--------------------------|------------------------------|
| (1) one mole of ammonia  | (2) two moles of nitric acid |
| (3) two moles of ammonia | (4) one mole of nitric acid  |

**Ans.** two moles of ammonia

116. Which one of the following ores is best concentrated by froth – floatation method?
- |               |                 |
|---------------|-----------------|
| (1) Magnetite | (2) Malachite   |
| (3) Galena    | (4) Cassiterite |

**Ans.** Galena

117. Beryllium and aluminium exhibit many properties which are similar. But the two elements differ in
- (1) exhibiting maximum covalency in compound
  - (2) exhibiting amphoteric nature in their oxides
  - (3) forming covalent halides
  - (4) forming polymeric hydrides

**Ans.** exhibiting maximum covalency in compound

118. Aluminium chloride exists as dimer,  $\text{Al}_2\text{Cl}_6$  in solid state as well as in solution of non-polar solvents such as benzene. When dissolved in water, it gives
- |                                     |   |
|-------------------------------------|---|
| (1) $\text{Al}^{3+} + 3\text{Cl}^-$ | (2) $\text{Al}_2\text{O}_3 + 6\text{HCl}$                   |
| (3) $[\text{Al}(\text{OH})_6]^{3-}$ | (D) $[\text{Al}(\text{H}_2\text{O})_6]^{3+} + 3\text{Cl}^-$ |

**Ans.**  $[\text{Al}(\text{H}_2\text{O})_6]^{3+} + 3\text{Cl}^-$

119. The soldiers of Napoleon army while at Alps during freezing winter suffered a serious problem as regards to the tin buttons of their uniforms. White metallic tin buttons got converted to grey powder. This transformation is related to
- (1) an interaction with nitrogen of the air at very low temperatures
  - (2) an interaction with water vapour contained in the humid air
  - (3) a change in the partial pressure of oxygen in the air
  - (4) a change in the crystalline structure of tin

**Ans.** a change in the crystalline structure of tin

120. The  $E^\circ_{M^{+3}/M^{2+}}$  values for Cr, Mn, Fe and Co are  $-0.41$ ,  $+1.57$ ,  $+0.77$  and  $+1.97$  V respectively. For which one of these metals the change in oxidation state from +2 to +3 is easiest?

- (1) Cr
- (2) Co
- (3) Fe
- (4) Mn

**Ans.** Cr

121. Excess of KI reacts with  $\text{CuSO}_4$  solution and then  $\text{Na}_2\text{S}_2\text{O}_3$  solution is added to it. Which of the statements is incorrect for this reaction?

- (1)  $\text{Cu}_2\text{I}_2$  is reduced
- (2) Evolved  $\text{I}_2$  is reduced
- (3)  $\text{Na}_2\text{S}_2\text{O}_3$  is oxidized
- (4)  $\text{CuI}_2$  is formed

**Ans.**  $\text{CuI}_2$  is formed

122. Among the properties (a) reducing (b) oxidising (c) complexing, the set of properties shown by  $\text{CN}^-$  ion towards metal species is

- (1) a, b
- (2) a, b, c
- (3) c, a
- (4) b, c

**Ans.** c, a

123. The coordination number of central metal atom in a complex is determined by

- (1) the number of ligands around a metal ion bonded by sigma bonds
- (2) the number of only anionic ligands bonded to the metal ion
- (3) the number of ligands around a metal ion bonded by sigma and pi- bonds both
- (4) the number of ligands around a metal ion bonded by pi-bonds

**Ans.** the number of ligands around a metal ion bonded by sigma

124. Which one of the following complexes is an outer orbital complex?

- (1)  $[\text{Fe}(\text{CN})_6]^{4-}$
- (2)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$
- (3)  $[\text{Co}(\text{NH}_3)_6]^{3+}$
- (4)  $[\text{Mn}(\text{CN})_6]^{4-}$

**Ans.**  $[\text{Ni}(\text{NH}_3)_6]^{2+}$

125. Coordination compounds have great importance in biological systems. In this context which of the following statements is incorrect?

- (1) Chlorophylls are green pigments in plants and contains calcium
- (2) Carboxypeptidase – A is an enzyme and contains zinc
- (3) Cyanocobalamin is  $\text{B}_{12}$  and contains cobalt
- (4) Haemoglobin is the red pigment of blood and contains iron

**Ans.** Chlorophylls are green pigments in plants and contains calcium

126. Cerium (Z = 58) is an important member of the lanthanoids. Which of the following statements about cerium is incorrect?

- (1) The common oxidation states of cerium are +3 and +4
- (2) Cerium (IV) acts as an oxidizing agent
- (3) The +4 oxidation state of cerium is not known in solutions
- (4) The +3 oxidation state of cerium is more stable than the +4 oxidation state

**Ans.** The +4 oxidation state of cerium is not known in solutions

127. Which one the following has largest number of isomers?

- (1)  $[\text{Ru}(\text{NH}_3)_4\text{Cl}_2]^+$
  - (2)  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$
  - (3)  $[\text{Ir}(\text{PR}_3)_2\text{H}(\text{CO})]^{2+}$
  - (4)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$
- (R = alkyl group, en = ethylenediamine)



**Ans.**  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$

128. The correct order of magnetic moments (spin only values in B.M.) among is

- (1)  $[\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-}$
  - (2)  $[\text{Fe}(\text{CN})_6]^{4-} > [\text{CoCl}_4]^{2-} > [\text{MnCl}_4]^{2-}$
  - (3)  $[\text{Fe}(\text{CN})_6]^{4-} > [\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-}$
  - (4)  $[\text{MnCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-} > [\text{CoCl}_4]^{2-}$
- (Atomic numbers: Mn = 25; Fe = 26, Co = 27)

**Ans.**  $[\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-}$

129. Consider the following nuclear reactions



The number of neutrons in the element L is

- (1) 142
- (2) 146
- (3) 140
- (4) 144

**Ans.** 144

130. The half – life of a radioisotope is four hours. If the initial mass of the isotope was 200 g, the mass remaining after 24 hours undecayed is

- (1) 1.042 g
- (2) 4.167 g
- (3) 3.125 g
- (4) 2.084 g

**Ans.** 3.125 g

131. The compound formed in the positive test for nitrogen with the Lassaigne solution of an organic compound is

- (1)  $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$
- (2)  $\text{Na}_4[\text{Fe}(\text{CN})_5\text{NOS}]$
- (3)  $\text{Fe}(\text{CN})_3$
- (4)  $\text{Na}_3[\text{Fe}(\text{CN})_6]$

**Ans.**  $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$

132. The ammonia evolved from the treatment of 0.30 g of an organic compound for the estimation of nitrogen was passed in 100 mL of 0.1 M sulphuric acid. The excess of acid required 20 mL of 0.5 M sodium hydroxide solution for complete neutralization. The organic compound is

- (1) acetamide
- (2) thiourea
- (3) urea
- (4) benzamide




**Ans.** urea

133. Which one of the following has the minimum boiling point?

- (1) n-butane (2) isobutane  
(3) 1-butene (4) 1-butyne

**Ans.** isobutane

134. The IUPAC name of the compound



- (1) 3,3-dimethyl-1-hydroxy cyclohexane (2) 1,1-dimethyl-3-cyclohexanol  
(3) 3,3-dimethyl-1-cyclohexanol (4) 1,1-dimethyl-3-hydroxy cyclohexane

**Ans.** 3,3-dimethyl-1-cyclohexanol

135. Which one of the following does not have  $sp^2$  hybridized carbon?

- (1) Acetone (2) Acetamide  
(3) Acetonitrile (4) Acetic acid


**Ans.** Acetonitrile

136. Which of the following will have meso-isomer also?

- (1) 2-chlorobutane (2) 2-hydroxypropanoic acid  
(3) 2,3-dichloropentane (4) 2,3-dichlorobutane

**Ans.** 2,3-dichlorobutane

137. Rate of the reaction



is fastest when Z is


- (1) Cl (2)  $\text{OCOCH}_3$   
(3)  $\text{OC}_2\text{H}_5$  (4)  $\text{NH}_2$

**Ans.** Cl

138. Amongst the following compound, the optically active alkane having lowest molecular mass is

- (1)
- (2)
- (3)
- (4)


**Ans.**




139. Consider the acidity of the carboxylic acids:  
 (1) PhCOOH (2) o-NO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>COOH  
 (3) p-NO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>COOH (4) m-NO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>COOH

**Ans.** o-NO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>COOH

140. Which of the following is the strongest base?



**Ans.**



141. Which base is present in RNA but not in DNA?  
 (1) Uracil (2) Thymine  
 (3) Guanine (4) Cytosine

**Ans.** Uracil

142. The compound formed on heating chlorobenzene with chloral in the presence concentrated sulphuric acid is  
 (1) gammexene (2) hexachloroethane  
 (3) Freon (4) DDT

**Ans.** DDT

143. On mixing ethyl acetate with aqueous sodium chloride, the composition of the resultant solution is  
 (1) CH<sub>3</sub>COOC<sub>2</sub>H<sub>5</sub> + NaCl (2) CH<sub>3</sub>Cl + C<sub>2</sub>H<sub>5</sub>COONa  
 (3) CH<sub>3</sub>COCl + C<sub>2</sub>H<sub>5</sub>OH + NaOH (4) CH<sub>3</sub>COONa + C<sub>2</sub>H<sub>5</sub>OH

**Ans.** CH<sub>3</sub>COOC<sub>2</sub>H<sub>5</sub> + NaCl

144. Acetyl bromide reacts with excess of CH<sub>3</sub>MgI followed by treatment with a saturated solution of NH<sub>4</sub>Cl given  
 (1) acetone (2) acetyl iodide  
 (3) 2-methyl -2- propanol (4) acetamide

**Ans.** 2-methyl -2- propanol

145. Which one of the following reduced with zinc and hydrochloric acid to give the corresponding hydrocarbon?  
 (1) Ethyl acetate (2) Butan -2-one  
 (3) Acetamide (4) Acetic acid





**Ans.** Butan -2-one

146. Which of the following undergoes reaction with 50% sodium hydroxide solution to give the corresponding alcohol and acid?


- (1) Phenol (2) Benzoic acid  
(3) Butanal (4) Benzaldehyde

**Ans.** Benzaldehyde

147. Among the following compound which can be dehydrated very easily is

- (1)  (2) 
- (3)  (D) 

**Ans.**



148. Which of the following compound is not chiral?

- (1) 1-chloropentane (2) 3-chloro-2-methyl pentane  
(3) 1-chloro-2-methyl pentane (4) 2-chloropentane

**Ans.** 1-chloropentane

149. Insulin production and its action in human body are responsible for the level of diabetes. This compound belongs to which of the following categories?

- (1) A co-enzyme (2) An antibiotic  
(3) An enzyme (4) A hormone

**Ans.** A hormone

150. The smog is essentially caused by the presence of

- (1)  $O_2$  and  $O_3$  (2)  $O_3$  and  $N_2$   
(3) Oxides of sulphur and nitrogen (4)  $O_2$  and  $N_2$

**Ans.** Oxides of sulphur and nitrogen